



Check Point	ONCOLOGISTS	PHARMACISTS	NURSES
	prescription phase	ordering & preparation phase	administration phase
Familiarise themselves with the ENHERTU [®] Summary of Product Characteristics (SmPC) which is available at www.medicines.org.uk	✓	✓	✓
Always refer to both ENHERTU [®] and trastuzumab deruxtecan	✓	✓	✓
Electronic systems <ul style="list-style-type: none"> • Check correct medication before clicking • Always select the correct medication in the electronic medical record • Ensure the medication prescribed is ENHERTU[®] (trastuzumab deruxtecan) and not trastuzumab or trastuzumab emtansine • Request use of brand names, where possible 	✓		
Do not abbreviate, truncate or omit any name on the prescription	✓		
Ensure the correct medication is clearly recorded in the prescription and patient notes	✓		✓
Check that protocols to avoid medication errors are in place at the hospital/site and that they are followed		✓	✓
Be aware when reading prescriptions that there are multiple types of medication with a similar INN (e.g. trastuzumab, trastuzumab SC, trastuzumab emtansine and trastuzumab deruxtecan)		✓	
Double check the intended medication is ENHERTU [®] (trastuzumab deruxtecan) and that both are entered in the prescription and/or medical history		✓	
In case of any doubt, consult with the treating physician		✓	✓
Familiarise yourself with the different cartons, labels and cap colours available for all trastuzumab containing products to select the correct carton		✓	✓
Check vials have the labels specific to ENHERTU [®]		✓	✓
Ensure the correct medication is ordered from the wholesaler and that the correct medication is received in the pharmacy		✓	
Store ENHERTU [®] in a different place in the fridge to other trastuzumab containing products (e.g. Herceptin [®] or Kadcyca [®])		✓	
On receipt of the infusion bag, check the label on the infusion bag against the prescription and patient notes			✓
Consider using a two nurse double-checking system prior to infusion to ensure that the appropriate product and dosage is administered			✓
Remember the maximum dose of ENHERTU [®] is 5.4 mg/kg once every 3 weeks	✓	✓	✓
Remember ENHERTU [®] is diluted with 5% glucose solution		✓	✓
Familiarise yourself with the ENHERTU [®] dose modification for toxicities	✓		✓